Early Childhood Brain Development



Todd Twogood, MD, FAAP
Pediatrician



NUTRITION



How it effects Brain Growth & Development



Doctors believe

It is their responsibility to be involved in promoting quality child care in their practice community


Healthy Childcare American Newsletter 1999

Early Brain Development

- All behavioral development has to do with the brain
- Brain development is dependent upon both experience and genetics

The Brain's Composition

- Growth of the brain occurs from the inside out and the bottom up
- You are born with 100 billion brain cells
- There are 15,000 synaptic connections for each cell



Early Experiences are Crucial




- By age 3, 80% of synaptic connections are already made
- By the second decade of life growth levels off and pruning begins
- Increased experiences define the wiring of an infant's brain



Plasticity


The brain's ability to change as a result of experience.




Human Brain at Birth


6 Years Old

14 Years Old









Biology & Environment
Work together to shape the brain.



Nature vs. Nurture

- During the first 10 years of life the brain is twice as active as that of an adult's
- 60% of nutrition is used by the brain during the first year of life. This decreases to 30% by age 3
- Genetic and environmental factors have a more dynamic, qualitative interplay that can not be reduced to a simple equation

Growing a Healthy Brain

- Nurturing experiences.
- Good nutrition.
- Intervening early.
- Protection.
- Taking care of the caregiver.



Early Care Experiences

- Babies thrive when they receive warm, responsive early care
- Early care has a decisive, long lasting impact on how people develop, their ability to learn, and their capacity to regulate their own emotions



Parenting

The parent/child relationship is the most powerful influence on children's early brain development, particularly in the first two years.

Problem-based play

Regular, consistent play provides

rich stimulation

that is absorbed into core brain development.



Making a Difference

Parents need support from others to do the best job they can.

They always have.

Percent of Mothers Working Outside the Home

- 76% of mothers with children ages 0-6
- 85% of mothers with children ages 6-12
- 61% of mother of infants and toddlers
- 82,000 children potentially need child care
- 30% of infants and toddlers are in child care full time

Quality Care for Children

- The way that parents and caregivers relate to young children and the way they mediate children's contact with the environment directly effects the formation of neural pathways
- The child care environment provides an incredible opportunity to positively effect child development



The Challenge of Health and Safety

- Staff knowledge and skill the number one indicator of quality
- The turnover rate of providers is 30%
- The average hourly wage for a child care center worker is \$7.20
- Infants and toddlers in center care are cared for by an average of seven caregivers a year

National Cost, Quality and Child Outcomes Study

- 8% of infant-toddler programs rated high enough to support children's development
- 52% of infant-toddler programs rated mediocre (healthy and safe, some nurturing, few learning experiences)
- 40% of infant-toddler programs judged of poor quality with basic health and safety, no warmth or nurturing, no learning is encouraged


Child Care Quality in ND

(90 centers participating impacting 800 infants and toddlers)

- Only 6% of infant-toddler programs rated high enough to support children's development (national 8%)
- 76% of infant-toddler programs rated mediocre
- 18% of infant-toddler programs judged of poor quality with basic health and safety needs not met and care damaging to children's development



Figure 7
Rates of return to human capital investment initially setting investment to be equal across all ages



Rates of return to human capital investment initially setting investment to be equal across all ages



Critical Periods in Development

Critical periods are sensitive periods when certain important psychological processes are developing most rapidly.

Disturbances during these periods might alter the development of these processes in critical ways.

Examples of Critical Periods

- •German Measles during the 1sttrimester of pregnancy = death, or deaf, blind, deaf and blind, and heart problems.
- •Chimpanzees reared in darkness the first year of life will have permanently impaired vision because of certain cells of the retina fail to develop. The first year of life is critical for most normal visual development.

Examples of Critical Periods

- •Extremely poor nutrition can affect the central nervous system and brain growth.
- •Children born with a cataract will become permanently blind in that eye of the cataract if it is not promptly removed.

Effects of Neglect on a Child's Development

Growth delay with certain postural and behavioral signs. Infantsmay also exhibit mental and motor delays.

- A. Less adaptive social interaction behavior
- B. Less positive affective behavior
- C. Avoidant and resistant attachment to primary caretaker Deficits on measures of intelligence
- Deficits in language ability (auditory comprehension and verbal ability)

Academic underachievement

See special Issue: (1993). Child Neglect. Crimnal Justice and Behavior, 20, 49-64.i

Brain Research and Neglect

- Infants of depressed mothers exhibit atypical frontal electrical brain activity
- Chronic and severe neglect can reduce the size of the cerebral cortex thus the neglected child's brain can be smaller
- Stress early in life can promote long-term changes in multiple neural transmitter systems and brain structures (demonstrate through animal studies)

Impact of Neglect on Infant Mental Health

During Infancy....

- Attachment disorders
- •Delayed development
- •Infant depression
- •Inconsolable crying

During Toddler Years....

- Aggressive behavior
- •Impulsive behavior
- •Difficulty being motivated to learn
- •Difficulty tuning into teaching

Greater Likelihood for....

- •Placement in special education
- •Grade retention
- •School Dropout

The Developing Brain

Brain development is much more vulnerable to environmental influence than we ever suspected. Poor nutrition before and after birth can seriously interfere with brain development.

Brain scan studies of animals show changes in brain structure and function as a result of early experience.

Studies of children reared in poor environments show cognitive deficits by 18 months of age.


The Developing Brain


The influence of early environment on brain development is long lasting


Moderately to severely neglected children are at risk of having smaller head sizes resulting in smaller brains and lower IQs. PET scans of institutionalized children show the effect of extreme deprivation.


Healthy Brain

This PET scan of the brain of a normal child shows regions of high (red) and low (blue and black) activity. At birth, only primitive structures such as the brain stem (center) are fully functional; in regions like the temporal lobes (top). early childhood experiences wire the circuits.









How the brain develops


- Development of the brain and nervous system of the embryo begins shortly after conception
- Ectoderm → neural tube → spinal cord & brain (2-4 weeks)
- Umbilical cord and placenta not formed until week 2 of the embryos life

How the brain develops


- Period of the fetus (3rd month)
- Nervous system (brain & spinal cord) connect with organs and muscles
- Weeks 13-24, brain neurons are mostly complete (2nd trimester)



Human Brain Development – Synapse Formation




Sensitive Periods for Early Development




The Developing Brain

- •The brain development that takes place before age one is more rapid and extensive then we previously realized.
- •Position Emission Tomography (PET) studies show that biochemical patterns of a one-year-old' sbrain resembles a normal young adult.

OFF THE MARK/ by Mark Parisi



The New Hork Times Magazine



• Prenatal:


- Inadequate nutrition
- Illegal drugs alcohol
- Exposure to toxins
- Prescription drugs
- O-T-C drugs
- Stress

Early Negative Impact

The Developing Brain Smoking and ADHD


Evidence of an association between maternal smoking during pregnancy and attention deficit hyperactivity disorder.


Mothers who smoked during pregnancy had children with lower IQs compared with those whose mothers did not smoke. Source: Journal of Clinical Child Psychology, 1998



"CAN YOU GIVE HIM SOMETHING FOR MY NERVES?







Disproportionate exposure of Children

- Pound for pound, children take in more air, food, and water than adults
- Gastrointestinal tract is more permeable to nutrients and contaminants, e.g. calcium, Lead
- Children's activities and environment differ from adults

Disproportionate susceptibility of Children

- Developing tissues generally more vulnerable to environmental factors
- Environmental factors can "program" genetic expression, tissue and organ development in unique and lifelong ways
- Brain, respiratory, gastrointestinal, reproductive, immune system, metabolic systems are not fully developed for days, weeks, months, or years



The Developing Brain Poverty and Nutrition

Monckebergstudied 500 preschool children in Chile who were divided into three groups:

Group A-well nourished, middleclass children Group B-disadvantaged who received supplements of milk and free medical assistance Group C-disadvantaged, no assistance.

Results:

Groups A & B had little or no malnutrition, only 3% of children had below average IQs.

Group C had high incidence of malnutrition and 40% had IQs below 80, they also had smaller head sizes.



Nutrition

Optimal early child development begins with adequate nutrition from conception onwards.

Nutrition & Brain Development





- Prenatal nutrition
- Rapid development early in life
- Importance of nutrition in the main stages of development





- Conception to 2nd week
 - Umbilical cord not formed
 - Mother's food intake does not have an effect on the embryo

- Week 2 to birth
 - Nutrients to unborn child via mother
 - Most critical time for development of central nervous system
 - Folic acid
 - Early in pregnancy
 - Also late in pregnancy

- Folic acid deficiency (Biotech Week, 2004)
 - Anencephaly =irregular brainformation
 - Stem cell division
 - Lifelong brain impairment



- Proper nutrition
- Recommended amount of nutrients
- Alcohol & smoking
- Normal cognitive development



Nutrition for Infants



- Breast Milk vs. Formula
- Convenience vs. health
- 3rd world vs. developed countries

Nutrition for Infants



- (Better Nutrition, 2000)
 - Premature infants
 - Long-chain polyunsaturated fatty acids
 - Faster brainstem maturation

Nutrition for Infants

- When the bottle may be better
 - (Lawson, 2003)
 - Vegan mother's
 - Vitamin B12 deficiency
 - Developmental delays
 - Attention span, notivation

Nutrition in young children

- (Core, 2003)
 - Poor nutrition = hampered mental development
 - Information processing
 - Language
 - Below average for age
 - Marasumus



Nutrition for Continued Growth

- Healthy mind
 - Balanced diet
- Essential nutrients
 - variety



Summary

- Most critical stages of cognitive development
- Effects of poor nutrition
- Healthy eating
- Healthy mind





The end!

