Pediatric Headaches

James Miles, MD NDAFP Family Medicine Update January 21, 2022

Improving Health, Enriching Life

Disclosures

Relevant Financial Relationship(s)
None

Off Label Usage None

*****Altru

-Altru

Learning Objectives

- Review the classification of pediatric headaches
- Appreciate when there is a need for further evaluation (red flags)
- Understand the differences between migraine presentation in children and adults
- Discuss migraine treatment, preventative and acute measures

*Altru

Classification: Primary

- Migraine
- Tension-type HA
- Cluster HAs

*Altru

Improving Health, Enriching Lit

4

Tension-Type Headaches

- · Last 30 minutes to 7 days
- · Bilateral location

3

- · Pressing or tightening
- Mild to moderate intensity
- Not aggravated by physical activity
- · No nausea or vomiting
- · No more than one of photo- or phonophobia
- Tx: similar to migraine

***Altru**

Cluster Headaches

- Severe unilateral orbital, supraorbital, and/or temporal pain lasting 15 to 180 minutes
- · Conjunctival injection and/or lacrimation
- · Nasal congestion and/or rhinorrhea
- · Forehead and facial sweating
- · Miosis, ptosis, and/or eyelid edema
- · Sense of restlessness or agitation
- Tx: 100% oxygen or sumatriptan 6 mg IM acutely, verapamil 240 mg daily preventative

mproving Health, Enriching Life

Classification: Secondary

- Acute febrile illness
 - Most common children
 - Recurrent rhinosinusitis most common misDx
- CNS infection
- Posttraumatic
- Hypertension

- Medication
- · Medication overuse
- · Brain tumor
- Hydrocephalus
- Intracranial hemorrhage
- Idiopathic Intracranial Hypertension



niproving ready, Environing En



c

Clinical Presentation: Younger Kids

- · Able to attenuate or ignore through play
- Cry, rock, hide
- Chronic pain associated with anxiety, depression, and behavior problems
- · May affect ability to eat, sleep, or play



Improving Health, Enriching Life

9

History

- Age of onset
 - Anything occur around time of onset?
 - Current triggers?
- · Timing, frequency, and pattern
- Location
- · Characterizing headaches
 - Dull and achy vs. throbbing
 - "Draw the headache"
- Duration
- Severity

*Altru

10

History

- Associated signs and symptoms
- Previous evaluations
- · Headache hygiene
 - Sleep
 - Water
 - Diet
 - Activity
 - Caffeine use

routna Health Enrichina I ffe



_ _ . . .

History

- · Current and previous medications
- Pregnancy, labor and delivery, neonatal period
- Growth and development
- Major surgeries or hospitalizations
- · Chronic illnesses or medications
- Academic performance
- Family History

*****Altru

Improving Health, Er

Red Flags

- Young age
- Awakens in middle of
 Recumbent position night
- Occipital HAs
- Persisting, worsening
 Chronic illness
- · "Worse HA of my life"
- New HA
- Valsalva maneuver
- · Resistant to treatment
- Abnormal neurologic exam



13

Physical Exam

- · Most sensitive indicator of needing further evaluation
- Vital signs
 - Temperature
 - Blood pressure
 - Height and weight
 - Head circumference



14

Physical Exam: Neuro

- · Mental status
- · Cranial nerves
- Motor
- Sensation
- Reflexes
- Coordination
- Gait

17

*****Altru


Diagnostics

- · MRI brain vs. CT head
 - Nonspecific T2 hyperintensities w/ migriane
- MRA vs. CTA
- · LP: intracranial infection, subarachonoid hemorrhage, IIH
- CBC, toxicology, thyroid function tests
- · EEG not indicated



16

15



Pathophysiology

- · Polygenetic and multifactiorial
- · Cortical spreading depression
- · Serotonin and central pain control pathways
- CGRP

18

*Altru

Epidemiology

- Prevalence 2.5% <7 yrs, 5% by age 10
- Females>males after puberty
- Lower socioeconomic status associated with chronic migraine
- Family history of migraine common
- 90% adults diagnosed with recurrent sinus HAs actually have migraine HAs



improving riedial, Ethiciling Li

19

Chronic Migraine

- At least 15 HA days per month for more than 3 months
- Migraine features at least 8 days per month



Improving Health, Enriching Life

21

Migraine Subtypes

- Migraine with Aura
 - Brainstem aura
 - Rare
 - Females>Males
 - 7-20 years of age
 - Vertigo, dysarthria, tinnitus, diplopia, bilateral visual symptoms, bilateral paresthesias, decreased LOC, hypacusis
 - No motor weakness



Diagnostic Criteria

- · At least 5 attacks
- Lasting 4-72 hrs (2-72 in children)
- At least 2 of the following:
 - Unilateral (bilateral in children)
 - Pulsating/throbbing
 - Moderate to severe pain
 - Aggravation or causing avoidance of physical activity
- · At least one of the following:
 - Nausea and/or vomiting
 - Photophobia and phonophobia (behavioral in kids)



20

Migraine Subtypes

- Migraine with Aura
 - 14-30% children report aura
 - Typical aura
 - · Visual, sensory, and/or speech/language
 - No motor weakness
 - · Gradual development
 - Duration no longer than 1 hour
 - · Complete reversibility
 - Hemiplegic migraine
 - Sporadic vs. familial



22

Migraine Subtypes

- Migraine with Aura
 - Vestibular migraine
 - Any age
 - Severe vertigo/dizziness
 - 5-72 hours



Improving Health, Enriching Life

23

Migraine Subtypes

- · Migraine with aura
 - Retinal/ocular migraine

 - · Sudden loss of vision, photopsia, or scintillations in only one eye
 - 5-60 minutes
 - May occur with or without HA, typically uni-ocular
 - · Permanent vision loss may occur
 - · Fundoscopy: retina pale, constricted vessels
- Menstrual migraine

25



Migraine Equivalents/Variants

- Cyclic vomiting syndrome
- · Abdominal migraine
- Benign paroxysmal vertigo
- · Benign paroxysmal torticollis
- · Colic?
- · Acute confusional migraine
- · Alice in Wonderland syndrome
- Opthalmoplegic migraine



26

Migraine Complications

- Status migrainosus: attack lasting >72 hrs
- · Persistent aura without infarction
- Migrainous infarction (neuroimaging)
- Migraine aura-triggered seizure



27

Migraine Treatment



*Altru

28

30

Lifestyle Modifications

Education

29

- Good sleep hygiene
- Adequate water intake
- · Routine, healthy meals and snacks
- Regular exercise
- Avoid migraine triggers
- · Limit caffeine intake



Preventative Medications

- Indications
 - Frequent or long lasting
 - Significant disability or diminished quality of
 - Contraindication, failure of, or adverse effects of acute therapies
 - Medication overuse HA
 - Menstrual migraine



Preventative Medications

- Duration of treatment
 - 6-8 weeks at therapeutic doses for full benefit
 - -6-12 months of good headache control before discontinuing therapy



Altru

31 32

Preventative Medications: Tricyclics

Preventative Medications

- Riboflavin (Migrelief: 0.5-1 tab BID)

- Melatonin (1-3 mg ghs, max 9 mg ghs)

· Bright yellow/orange urine · GI upset uncommon

ginkolide B, polyunsaturated fats

· Daytime sleepiness

CHAMP trial: placebo is as effective and better

- Magnesium, feverfew, coenzyme Q10, butterbur,

tolerated than amitriptyline or topiramate

· Amitriptyline

Nutraceuticals

- 5-12.5 mg qhs, max 2 mg/kg/day or 100 mg qhs
- Sedating
- Tachychardia
- Prolongation of QT interval
 - EKG at baseline and higher doses
- Nortriptyline
 - Less sedating
- Trazodone

34



-Altru

33

Preventative Medications

- 2-4 mg qhs; max 12 mg per day, divided BID

Cyproheptadine

- Liquid or tablet

Somnolence

- Appetite stimulation

Preventative Medications: Antiseizure Medications

- Topiramate
 - 1-2 mg/kg/day, max 50 mg BID
 - Weight loss, cognitive impairment, paresthesias, closed-angle glaucoma, hypohydrosis, nephrolithiasis
 - Teratogenicity and LBW
 - Only FDA approved migraine preventative medication for 12-17 years of age



Preventative Medications: Antiseizure Medications

- Valproate
 - 10-15 mg/kg/day divided BID
 - Weight gain, hepatotoxicity (<2 yrs), thombocytopenia
 - Teratogenic
 - CBC, AST, ALT
- Gabapentin
 - TID dosing
 - Restless leg syndrome or neuropathic pain
- Levetiracetam

36

Altru

Preventative Medications: Beta Blockers

- Propranolol
 - 1 mg/kg/day divided TID (multiple doses per day)
 - Abdominal migraines
 - Hypotension, bradycardia, emotional disturbances, nightmares
 - Monitor HR and orthostatic BP (tall/skinny/active patients)
 - Use caution in patients with asthma, diabetes, depression



improving riesian, Etiniching E

37

Preventative Medications: Calcium Channel Blockers

- Cinnarizine
 - Mild drowsiness and uncommon weight gain
- Flunarizine
 - Sturge-Weber syndrome
 - Sedation and weight gain
- · Not readily available in US



Improving Health, Enriching Life

39

Neurostimulation

· Trancutaneous nerve stimulation

Preventative Medications:

- Beta-1 selective blocker, can be safer than

nonselective beta blockers (propranolol)

Beta Blockers

• Propranolol LA

Atenolol

- Cefaly and Nerivio devices
- Can be used for preventative and acute treatment
- Drowsiness, well tolerated
- Costs \$500 (Cefaly) \$99 (Nerivio), insurance may not cover
- Transcranial magnetic stimulation
 - 0.9 Tesla magnetic field in a brief pulse over occipital area
 - Preventative and acute treatment
 - Rented for \$750 every 3 months



₹Altru

40

Botulinum toxin

- Shown to decrease number of headache days in adults with chronic migraine
- Retrospective studies in children show similar results, clinical trial recently showed no benefit
- · IM injections every 12 weeks
- Neck/musculoskeletal pain, facial drooping
- · Must fail 2-3 prescription medications
- Yearly cost \$14K, typically well covered by insurance



Improving Health, Errich

CGRP Antagonists

- CGRP: found in unmyelinated sensory nerve fibers
 - Associated with transmission of painful stimuli
- Monoclonal antibodies to CGRP or its receptor have been shown to be effective in adults
- Self injection once per month
- Not approved for <18 years
 - Poorly covered by insurance, yearly cost >\$7K
 - Questionable efficacy

*****Altru

Other Preventative Treatments

- · Behavioral therapy
 - Cognitive behavioral therapy
 - Clinical trial showed benefit in children with CBT+amitriptyline vs. HA education+amitriptyline
 - · Comorbid anxiety and depression VERY common
 - Biofeedback
 - Limitations: availability, cost, patient ability and compliance



Improving Health, Enriching Life

43

Other Preventative Treatments

- Physical therapy
 - Chronic migraine
 - Head injury/concussion
 - Other comorbidities
- Other
 - Chiropractics
 - Massage therapy
 - Acupuncture/acupressure
 - Daith piercing

Improving Health, Enriching Life

- Marijuana and CBD



44

Menstrual Migraine

- · Hormonal therapy
- · "Mini-prophylaxis"
 - 1-2 days prior to expected headache onset
 - Continued for expected duration
 - Naproxen 550 mg BID
 - Long acting triptans



Improving Health, Enriching Life

45

Acute Treatment

- General measures
 - Dark, quiet room
 - Cool cloth applied to forehead
- NEVER USE:
 - Opioids
 - Barbituates
 - Benzodiazepines



Improving Health, Enrichin

46

Acute Treatment

- Analgesics
 - Ibuprofen 10 mg/kg or acetaminophen 15 mg/kg
 - Onset of migraine symptoms, can repeat in 2
 bours
 - Naproxen 5 mg/kg every 8-12 hours
 - Excedrin Migraine 1 tablet
 - · Adolescents only due to aspirin
 - No more than 2 doses per day, 2 days per week
 - Can take with caffeine.



(ltrii)

Acute Treatment

- Triptans
 - 5 years of age and older
 - Refractory to analgesics
 - Onset of migraine symptoms, can repeat in 2 hours
 - No more than 2 doses per day, 2 days per week
 - Can take with caffeine



Improving Health, Enriching

48

7

Acute Treatment

- Triptans
 - Sumatriptan 25-50 mg oral, 5-10 mg nasal
 - Rizatriptan 2.5-5 mg oral or dissolvable
 - Zolmitriptan 2.5-5 mg oral, dissolvable, or nasal
- · Nasal sprays: tilt head forward, hard candy
- · Contraindications: Hx ischemic vascular disease or arrhythmias a/w accessory conduction pathway d/o's
- Caution: brainstem aura and hemiplegic migraine

Acute Treatment

- Ditans
 - Lasmiditan
 - Target and activate 5-HT_{1F} receptors
 - Currently Phase I trials for children
 - Concern for increased risk for cardiovascular (CV) disease and CV events, especially in children



50

49

Acute Treatment

- Combination
 - Same time: triptan + naproxen
 - Severity based: analgesic mild to mod, triptan severe
- Antiemetics
 - Promethazine 0.25-0.5 mg/kg per dose
 - Zofran 4 mg

Altru

*****Altru

Status Migrainosus

- Clinic/Urgent Care
 - Ketorolac 15-60 mg IM + antiemetic
- · Emergency setting
 - IV fluids 20 mL/kg NS, max 1 L
 - IV ketorolac
 - IV antiemetic (ex. Prochlorperazine)
- · DHE: caution with brainstem aura and hemiplegic migraine
- · IV valproate

*Altru

52

Prognosis

- · Many patients improve over time
 - Rule of thirds
- · Girls more likely to relapse
- Early age may mean a less favorable prognosis



Resources

- Abu-Arafeh I, Razak S, Sivaraman B, Graham C. Prevalence of headache and migraine in children and adolescents: a systematic review of population-based studies. Dev Med Child Neurol 2010; 52:1088.
- El-Chammas K, Keyes J, Thompson N, et al. Pharmacologic treatment of pediatric headaches: a meta-analysis. JAMA Pediatr 2013; 167:250.

 Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. lalgia 2018; 38:1.
- Oskoui M, Pringsheim T, Holler-Managan Y, et al. Practice guideline update summary: Acute treatment of migraine in children and adolescents: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology and the American Headache Society. Neurology 2019; 93:487.



54

53

Resources

- Powers SW, Coffey CS, Chamberlin LA, et al. Trial of Amitriptyline, Topiramate, and Placebo for Pediatric Migraine. N Engl J Med 2017; 376:115.
- Silberstein SD. Practice parameter: evidence-based guidelines for migraine headache (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology 2000; 55:754.
- US Headache Consortium. Evidence-based guidelines in the primary care setting: Neuroimaging in patients with nonacute headache, 2000. www.aan.com/professionals/practice/pdfs/ql0088.pdf
- UpToDate. https://www.uptodate.com



Improving Health, Enriching Life

